

Genus *Liptenara* Bethune-Baker, 1915

Annals and Magazine of Natural History (8) **16**: 186 (186-203).

Type-species: *Liptenara batesi* Bethune-Baker, by original designation.

A purely Afrotropical genus containing three species.

**Liptenara batesi* Bethune-Baker, 1915

Liptenara batesi Bethune-Baker, 1915. *Annals and Magazine of Natural History* (8) **16**: 187 (186-203).

Liptenara batesi Bethune-Baker, 1915. d'Abbrera, 2009: 613.

Type locality: Cameroon: "Bitje, Cameroons".

Distribution: Cameroon, Gabon, Democratic Republic of Congo (Equateur, Tshuapa, Tshopo).

Specific localities:

Cameroon – Bitje (TL).

Gabon – Tchimbele (van de Weghe, 2010); Kebiri Forest, Bateke Plateau (van de Weghe, 2010).

Early stages: Nothing published.

Larval food: Nothing published.

**Liptenara hiendlmayri* (Dewitz, 1887)

Pentila hiendlmayri Dewitz, 1887. *Deutsche Entomologische Zeitschrift* **30**: 429 (427-430).

Liptenara hiendlmayri Dewitz, 1887. d'Abbrera, 2009: 613.

Type locality: [Democratic Republic of Congo]: "Mukenge".

Distribution: Congo, Democratic Republic of Congo (Lulua, Sankuru), Uganda (west).

Specific localities:

Democratic Republic of Congo – Mukenge (TL).

Uganda – Bwamba; Budongo Forest; Semuliki N.P. (Davenport & Howard, 1996).

Early stages: Nothing published.

Larval food: Nothing published.

**Liptenara schoutedeni* (Hawker-Smith, 1926)

Pentila schoutedeni Hawker-Smith, 1926. *Revue Zoologique Africaine* **14**: 238 (237-241).

Liptenara schoutedeni Hawker-Smith, 1926. d'Abbrera, 2009: 613.

Type locality: [Democratic Republic of Congo]: "Sandoa; Kayombo".

Distribution: Democratic Republic of Congo (south-east – Lualaba).

Specific localities:

Democratic Republic of Congo – Sandoa (TL); Kayombo (Hawker-Smith, 1926).

Early stages: Nothing published.

Larval food: Nothing published.